

Prospects for Discovery of DM Annihilation to Primary Neutrinos with IceCube

Bradley Knockel
University of New Mexico

Working with...
Rouzbeh Allahverdi (UNM), Alexander Friedland (LANL),
Katherine Richardson (UNM)

INFO 13
Santa Fe, USA, Aug 26-30

Outline


(1) Introduction

(2) Methods and Results

(3) Conclusions

What's the matter?

Our Present Universe



$$\Omega_0 = 0.683 + 0.268 + 0.049 = 1$$

The WIMP Explanation

- WIMPs (**Weakly Interacting Massive** Particles) are currently the best explanation for the composition of dark matter
- Particles are most likely **weakly interacting**. Not too strong, but we assume stronger than gravity
- Particles are **massive** (that is, not massless) to explain observed structure formation

WIMP Relic Density via Thermal Freeze-Out

$$\Gamma_{\text{ann}} \sim H$$






Fig. 5.1: The freeze out of a massive particle species. The dashed line is the actual abundance, and the solid line is the equilibrium abundance.

$$\langle\sigma_{\text{ann}}v\rangle = 3 \times 10^{-26} \frac{\text{cm}^3}{\text{s}}$$

IceCube




ice cube



Ice Cube



Indirect Detection via the Sun



$$\Gamma_{\text{ann}} = \frac{\Gamma_{\text{cap}}}{2} \tanh^2\left(\frac{t}{\tau_{\text{eq}}}\right) \approx \frac{\Gamma_{\text{cap}}}{2}$$

and B_i

Why Prompt Neutrinos

- The LHC has constrained the MSSM, so the neutralino may not be the WIMP
- Our analysis is model independent, and we are not biased towards the neutralino
- Many models have enhanced prompt neutrinos
- For example, if the WIMP has lepton number, it can annihilate to $\nu\nu$

Enhanced prompt neutrinos

Lindner, Merle, and Niro, Phys. Rev. D 82, 123529 (2010).

Annihilation Channel Examples


$$\chi + \chi \rightarrow \tau^- + \tau^+$$

$$\chi + \chi \rightarrow W^+ + W^-$$


$$\chi + \chi \rightarrow \nu_e + \bar{\nu}_e$$

$$\chi + \chi \rightarrow \nu_\mu + \bar{\nu}_\mu$$

$$\chi + \chi \rightarrow \nu_\tau + \bar{\nu}_\tau$$



Muon Neutrinos Reaching Detector



There are corresponding angular distributions as well.

Outline

(1) Introduction

(2) Methods and Results

(3) Conclusions

Our Analysis

(1) Acquire simulated data for signal and background at IceCube using DarkSUSY. We acquire “contained” and “through-going” muons for a low-energy and high-energy analysis, respectively.

(2) Model IceCube’s detection of the muon tracks using published effective detector volumes and areas and by smearing the angular and energy distributions by 1° and 40 GeV respectively


(3) Optimize the analysis for dark matter discovery at 90% confidence using a χ^2 analysis

$$\frac{S}{\sqrt{S+1.2B}} = 1.64$$

(4) Distinguish between different annihilation channels assuming a discovery and an independent measurement of mass

$$\sigma = \frac{|S_c - S_t|}{\sqrt{S_c + 1.2B_c}}$$

Benchmarking Using the W Channel




Effective Volume/Area of IceCube
Astropart. Phys. 35, 615 (2012).

IceCube 2011
Phys. Rev. D 85, 042002 (2012).

IceCube 2012
Physical Review Letters 110, 131302 (2013).

IceCube can be modeled simply.


Sensitivity Plot for All Channels



COUPP
Phys. Rev. D 86, 052001 (2012).

Factor of ~ 8 between ν and W
Allahverdi and Richardson, Phys. Rev. D 85, 113012 (2012).


Corresponding $\langle \sigma_{\text{ann}} v \rangle$ Required for Equilibrium in Sun



Sun might be the only way the detect prompt neutrinos.

Compare to galactic prompt-neutrino bounds of $\sim 10^{-23}$
IceCube (2011), e-Print: arXiv:1111.2738 [astro-ph.HE].

Distinguishing Channels (asking if ν_μ fits the data)



Highly sensitive to the energy smearing that we choose to model IceCube

Outline

(1) Introduction

(2) Methods and Results

(3) Conclusions


Conclusions

- We should consider prompt neutrinos
- Although the solar community ignores prompt neutrinos, indirect detection via the Sun may be the only way to detect them
- IceCube is sensitive to prompt neutrinos for DM mass < 1 TeV
- Prompt neutrinos can be distinguished from other annihilation channels


Paper on arXiv soon!

Additional Slides


Sensitivity (τ and ν_μ mixtures)




mixtures of τ and ν_μ




Distinguishing using 10 GeV smearing instead of 40 GeV (for low-energy)



τ as target



Discovery Cuts



First energy bin begins ABOVE energy cut,
and angle is integrated up to the angular cut.

Distinguishing Angular Cuts

